

全国订货电话 4008-723-722

兔抗 ADD2 多克隆抗体

- 中文名称: 兔抗 ADD2 多克隆抗体
- 英文名称: Anti-ADD2 rabbit polyclonal antibody
- 别 名: ADDB
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 抗 原: ADD2
- 宿 主: Rabbit
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格



| | Adducins are heteromeric proteins composed of differ |
|-----------------------------|---|
| Background: | ent subunits referred to as adducin alpha, beta and |
| | gamma. The three subunits are encoded by distinct g |
| | enes and belong to a family of membrane skeletal pr |
| | oteins involved in the assembly of spectrin-actin netw |
| | ork in erythrocytes and at sites of cell-cell contact in |
| | epithelial tissues. While adducins alpha and gamma a |
| | re ubiquitously expressed, the expression of adducin |
| | beta is restricted to brain and hematopoietic tissues. |
| | Adducin, originally purified from human erythrocytes, |
| | was found to be a heterodimer of adducins alpha an |
| | d beta. Polymorphisms resulting in amino acid substit |
| | utions in these two subunits have been associated wi |
| | th the regulation of blood pressure in an animal mo |
| | del of hypertension. Heterodimers consisting of alpha |
| | and gamma subunits have also been described. Struc |
| | turally, each subunit is comprised of two distinct do |
| | mains. The amino-terminal region is protease resistan |
| | t and globular in shape, while the carboxy-terminal r |
| | egion is protease sensitive. The latter contains multipl |
| | e phosphorylation sites for protein kinase C, the bind |
| | ing site for calmodulin, and is required for associatio |
| | n with spectrin and actin. Alternatively spliced transcri |
| | pt variants have been described. |
| Applications: | ELISA, IHC |
| Name of antibody: | ADD2 |
| Immunogen: | Fusion protein of human ADD2 |
| Full name: | adducin 2 (beta) |
| Synonyms : | ADDB |
| SwissProt: | P35612 |
| ELISA Recommended dilution: | 1000-2000 |
| IHC positive control: | Human brain |
| IHC Recommend dilution: | |



全国订货电话 4008-723-722

