

兔抗 ATXN7L3 多克隆抗体

- 中文名称:兔抗 ATXN7L3 多克隆抗体
- 英文名称: Anti-ATXN7L3 rabbit polyclonal antibody
- 相关类别: 一抗
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗原: ATXN7L3
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	Component of the transcription regulatory histone a cetylation (HAT) complex SAGA, a multiprotein com plex that activates transcription by remodeling chro matin and mediating histone acetylation and deubiq uitination. Within the SAGA complex, participates in a subcomplex that specifically deubiquitinates both histones H2A and H2B. The SAGA complex is recruit ed to specific gene promoters by activators such as MYC, where it is required for transcription. Required for nuclear receptor-mediated transactivation. Within the complex, it is required to recruit USP22 and EN Y2 into the SAGA complex.
Applications:	ELISA, WB, IHC
Name of antibody:	ATXN7L3
Immunogen:	Fusion protein of human ATXN7L3





Full name:	ataxin 7-like 3
SwissProt:	Q14CW9
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human brain and human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	39 kDa
WB Positive control:	Hela and Raji cell lysates
WB Recommended dilution:	500-2000





