

兔抗 C7 多克隆抗体

中文名称：兔抗 C7 多克隆抗体

英文名称：Anti-C7 rabbit polyclonal antibody

别名：complement component 7;

相关类别：一抗

抗原：C7

储存：冷冻（-20℃）

宿主：Rabbit

克隆类型：rabbit polyclonal

反应种属：Human

标记物：Unconjugate

技术规格

Background:

The complement cascade is a multi-protein system that functions to clear pathogens from an infected host. Part of the innate (unchanging) immune system, the complement cascade consists of proteins and inactive zymogens that are present in blood and are stimulated by one of several triggers. Once stimulated, the cascade relays amplified responses throughout the body, ultimately activating the cell-killing membrane attack complex which can insert itself into the cell membrane and cause the cell to lyse. C7 (complement component 7) is an 843 amino acid secreted protein that participates in the formation of membrane attack complex (MAC), a complex that forms pores in the plasma membra

	ne of target cells for innate and adaptive immune responses. As a membrane anchor, C7 exists as a monomer or dimer and can form multimeric rosettes with C5 β . C7 defects are the cause of component C7 deficiency (C7D), characterized by recurrent bacterial infections caused by <i>Neisseria meningitidis</i> .
Applications:	ELISA, WB
Name of antibody:	C7
Immunogen:	Fusion protein of human C7
Full name:	complement component 7
SwissProt:	P10643
ELISA Recommended dilution:	5000-10000
WB Predicted band size:	94 kDa
WB Positive control:	HUVEC cell lysate
WB Recommended dilution:	500-2000

