

兔抗 HDAC4 多克隆抗体

中文名称： 兔抗 HDAC4 多克隆抗体

英文名称： Anti-HDAC4 rabbit polyclonal antibody

别 名： HD4; AHO3; BDMR; HDACA; HA6116; HDAC-4; HDAC-A

相关类别： 一抗

储 存： 冷冻 (-20°C)

宿 主： Rabbit

抗 原： HDAC4

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

Background:	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.
Applications:	ELISA, IHC
Name of antibody:	HDAC4
Immunogen:	Synthetic peptide of human HDAC4
Full name:	histone deacetylase 4

Synonyms :	HD4; AHO3; BDMR; HDACA; HA6116; HDAC-4; HDAC-A
SwissProt:	P56524
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	25-100

