

兔抗 ICAM1 多克隆抗体

- 中文名称: 兔抗 ICAM1 多克隆抗体
- 英文名称: Anti-ICAM1 rabbit polyclonal antibody
- 别名: BB2; CD54; P3.58
- 相关类别: 一抗
- 抗原: ICAM1
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a cell surface glycoprotein which is typica lly expressed on endothelial cells and cells of the immune s ystem. It binds to integrins of type CD11a / CD18, or CD11 b / CD18 and is also exploited by Rhinovirus as a receptor. ICAM proteins are ligands for the leukocyte adhesion protei n LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-end othelial migration, ICAM1 engagement promotes the assembl y of endothelial apical cups through ARHGEF26/SGEF and R HOG activation. In case of rhinovirus infection acts as a cellu lar receptor for the virus.
Applications:	ELISA, WB, IHC
Name of antibody:	ICAM1
Immunogen:	Synthetic peptide of human ICAM1





Full name:	intercellular adhesion molecule 1
Synonyms :	BB2; CD54; P3.58
SwissProt:	P05362
ELISA Recommended diluti on:	2000-5000
IHC positive control:	Human lung cancer and Human breast cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	58 kDa
WB Positive control:	Huvec, hela and raji cells
WB Recommended dilution :	500-2000





全国订货电话 4008-723-722

