

小鼠抗 SUPT16H 单克隆抗体

中文名称： 小鼠抗 SUPT16H 单克隆抗体

英文名称： Anti-SUPT16H mouse monoclonal antibody

别 名： SPT16 homolog, facilitates chromatin remodeling subunit; CDC68; SPT16; FACTP140; SPT16/CDC68

相关类别： 一抗

储 存： 冷冻 (-20°C)

宿 主： Mouse

抗 原： SUPT16H

反应种属： Human

标 记 物： Unconjugate

克隆类型： mouse monoclonal

技术规格

Background:	Transcription of protein-coding genes can be reconstituted on naked DNA with only the general transcription factors and RNA polymerase II. However, this minimal system cannot transcribe DNA packaged into chromatin, indicating that accessory factors may facilitate access to DNA. One such factor, FACT (facilitates chromatin transcription), interacts specifically with histones H2A/H2B to effect nucleosome disassembly and transcription elongation. FACT is composed of an 80 kDa subunit and a 140 kDa subunit; this gene encodes the 140 kDa subunit.
Applications:	WB, IHC
Name of antibody:	SUPT16H

Immunogen:	Fusion protein of human SUPT16H
Full name:	SPT16 homolog, facilitates chromatin remodeling subunit
Synonyms:	CDC68; SPT16; FACTP140; SPT16/CDC68
SwissProt:	Q9Y5B9
IHC positive control:	Human pancreas tissue
IHC Recommend dilution:	50-150
WB Predicted band size:	120 KD
WB Positive control:	Jurkat, Hela cell lysates
WB Recommended dilution :	500-2000