

小鼠抗 TAPBPL 单克隆抗体

- 中文名称: 小鼠抗 TAPBPL 单克隆抗体
- 英文名称: Anti-TAPBPL mouse monoclonal antibody
- 别 名: TAPBP-R; TAPBPR
- 储 存:冷冻(-20℃) 避光
- 抗 原: TAPBPL
- 宿 主: Mouse
- 反应种属: Human, Dog, Rat, Monkey, Mouse
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: mouse monoclonal

技术规格

Background:	Tapasin, or TAPBP (MIM 601962), is a member of the variable-c onstant Ig superfamily that links major histocompatibility compl ex (MHC) class I molecules to the transporter associated with a ntigen processing (TAP; see MIM 170260) in the endoplasmic r eticulum (ER). The TAPBP gene is located near the MHC compl ex on chromosome 6p21.3. TAPBPL is a member of the Ig sup erfamily that is localized on chromosome 12p13.3, a region so mewhat paralogous to the MHC.[supplied by OMIM, Mar 2008].
Applications:	WB, IHC, IF
Name of antibody:	TAPBPL
Immunogen:	Fusion protein of human TAPBPL
Full name:	TAP binding protein-like (TAPBPL)
Synonyms:	TAPBP-R; TAPBPR





SwissProt:	Q9BX59
IHC positive control:	human pancreas tissue
IHC Recommend dilutio n:	30-150
WB Predicted band size:	50 kDa
WB Positive control:	HepG2, Hela, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 ce II lysates
WB Recommended diluti on:	200-1000