

## 小鼠抗 PIK3CG 单克隆抗体

- 中文名称: 小鼠抗 PIK3CG 单克隆抗体
- 英文名称: Anti-PIK3CG mouse monoclonal antibody
- 别 名: p110gamma; p120-PI3K; PI3CG; PI3K; PI3Kgamma; PIK3
- 储 存:冷冻(-20℃) 避光
- 抗 原: PIK3CG
- 宿 主: Mouse
- 反应种属: Human, Monkey, Mouse
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: mouse monoclonal

## 技术规格

Background:	This gene encodes a protein that belongs to the pi3/pi4-kinas e family of proteins. The gene product is an enzyme that phos phorylates phosphoinositides on the 3-hydroxyl group of the i nositol ring. It is an important modulator of extracellular signal s, including those elicited by E-cadherin-mediated cell-cell adh esion, which plays an important role in maintenance of the str uctural and functional integrity of epithelia. In addition to its r ole in promoting assembly of adherens junctions, the protein i s thought to play a pivotal role in the regulation of cytotoxicit y in NK cells. The gene is located in a commonly deleted seg ment of chromosome 7 previously identified in myeloid leuke mias. [provided by RefSeq, Jul 2008].
Applications:	WB, IHC, IF





	1000 120 122
Name of antibody:	PIK3CG
Immunogen:	Fusion protein of human PIK3CG
Full name:	phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit gamma (PIK3CG), transcript variant 1
Synonyms:	p110gamma; p120-PI3K; PI3CG; PI3K; PI3Kgamma; PIK3
SwissProt:	P48736
IHC positive control:	adenocarcinoma of human ovary tissue and human pancreas ti ssue; human kidney tissue and human liver tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	126 kDa
WB Positive control:	SVT2, COS7 cell lysates
WB Recommended diluti on:	200-1000