

兔抗 CNTNAP2 多克隆抗体

- 中文名称: 兔抗 CNTNAP2 多克隆抗体
- 英文名称: Anti-CNTNAP2 rabbit polyclonal antibody
- 别 名: contactin associated protein-like 2; CDFE; NRXN4; AUTS15; CASPR2; PTHSL1
- 抗 原: CNTNAP2
- 相关类别: 一抗
- 储 存: 冷冻(-20℃)
- 宿 主: Rabbit
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a member of the neurexin family wh ich functions in the vertebrate nervous system as cell a dhesion molecules and receptors. This protein, like other neurexin proteins, contains epidermal growth factor repe ats and laminin G domains. In addition, it includes an F 5/8 type C domain, discoidin/neuropilin- and fibrinogen- like domains, thrombospondin N-terminal-like domains a nd a putative PDZ binding site. This protein is localized at the juxtaparanodes of myelinated axons, and mediate s interactions between neurons and glia during nervous
	s interactions between neurons and glia during nervous system development and is also involved in localization
	of potassium channels within differentiating axons. This gene encompasses almost 1.5% of chromosome 7 and i

TW-reagent.com

全国订货电话 4008-723-722

	s one of the largest genes in the human genome. It is directly bound and regulated by forkhead box protein P 2 (FOXP2), a transcription factor related to speech and I anguage development. This gene has been implicated in multiple neurodevelopmental disorders, including Gilles de la Tourette syndrome, schizophrenia, epilepsy, autism , ADHD and mental retardation.
Applications:	ELISA, WB, IHC
Name of antibody:	CNTNAP2
Immunogen:	Fusion protein of human CNTNAP2
Full name:	contactin associated protein-like 2
Synonyms:	CDFE; NRXN4; AUTS15; CASPR2; PTHSL1
SwissProt:	Q9UHC6
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer;Human tonsil
IHC Recommend dilution:	20-100
WB Predicted band size:	148 kDa
WB Positive control:	Rat brain tissue , Mouse brain tissue and Human cerebr um tissue lysates
WB Recommended dilution:	200-1000





