

兔抗 CYP3A7 多克隆抗体

中文名称: 兔抗 CYP3A7 多克隆抗体

英文名称: Anti-CYP3A7 rabbit polyclonal antibody

别名: CP37; CYP3A7; P450-HFLA; P-450111A7; P-450(HFL33)

相关类别: 一抗

储存: 冷冻 (-20℃) 避光

宿主: Rabbit

抗原: CYP3A7

反应种属: Human

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	This gene encodes a member of the cytochrome P450 superfamily of enzymes, which participate in drug metabolism and the synthesis of cholesterol, steroids and other lipids. This enzyme hydroxylates testosterone and dehydroepiandrosterone 3-sulphate, which is involved in the formation of estradiol during pregnancy. This gene is part of a cluster of related genes on chromosome 7q21.1. Naturally-occurring readthrough transcription occurs between this gene and the downstream CYP3A7 pseudogene and is represented by GeneID:100861540.
Applications:	WB
Name of antibody:	CYP3A7
Immunogen:	Synthesized peptide derived from internal of human Cytochr

	ome P450 3A7.
Full name:	cytochrome P450, family 3, subfamily A, polypeptide 7
Synonyms :	CP37; CYP3A7; P450-HFLA; P-450111A7; P-450(HFL33)
SwissProt:	A4D288/P24462
WB Predicted band size:	58 kDa
WB Positive control:	LOVO cells lysate
WB Recommended dilution:	500-3000

