

## TRAF3IP2 抗原(重组蛋白)

中文名称: TRAF3IP2 抗原(重组蛋白)

英文名称: TRAF3IP2 Antigen (Recombinant Protein)

别名: TRAF3 interacting protein 2; ACT1; CIKS; C6orf2; C6orf4; C6orf5; C6orf6; PSORS13

- 储存: 冷冻(-20℃)
- 相关类别: 抗原
- 概 述:

Fusion protein corresponding to a region derived from 1-256 amino acids of human TRAF3IP2

技术规格:

| Full name:         | TRAF3 interacting protein 2  |
|--------------------|--|
| Synonyms:          | ACT1; CIKS; C6orf2; C6orf4; C6orf5; C6orf6; PSORS13  |
| Swissprot:         | O43734   |
| Gene Accession:    | BC002823   |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PAGE   |
| Expression system: | Escherichia coli   |
| Tags:              | His tag C-Terminus, GST tag N-Terminus   |
| Background:        | This gene encodes a protein involved in regulating responses to cytoki<br>nes by members of the Rel/NF-kappaB transcription factor family. Thes<br>e factors play a central role in innate immunity in response to pathog<br>ens, inflammatory signals and stress. This gene product interacts with<br>TRAF proteins (tumor necrosis factor receptor-associated factors) and e<br>ither I-kappaB kinase or MAP kinase to activate either NF-kappaB or J<br>un kinase. Several alternative transcripts encoding different isoforms ha<br>ve been identified. Another transcript, which does not encode a protei<br>n and is transcribed in the opposite orientation, has been identified. O |





verexpression of this transcript has been shown to reduce expression of at least one of the protein encoding transcripts, suggesting it has a regulatory role in the expression of this gene.