

兔抗 DMAP1 多克隆抗体

中文名称：兔抗 DMAP1 多克隆抗体

英文名称：Anti-DMAP1 rabbit polyclonal antibody

别名：EPD; PDE; ATQ1

相关类别：一抗

储存：冷冻（-20℃）

抗原：DMAP1

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a subunit of several, distinct complexes involved in the repression or activation of transcription. The encoded protein can independently repress transcription and is targeted to replication foci throughout S phase by interacting directly with the N-terminus of DNA methyltransferase 1. During late S phase, histone deacetylase 2 is added to this complex, providing a means to deacetylate histones in transcriptionally inactive heterochromatin following replication. The encoded protein is also a component of the nucleosome acetyltransferase of H4 complex and interacts with the transcriptional corepressor tumor susceptibility gene 101 and the pro-apoptotic death-associated protein 6,

	among others. Alternatively spliced transcript variants encoding the same protein have been described.
Applications:	ELISA, WB, IHC
Name of antibody:	DMAP1
Immunogen:	Synthetic peptide of human DMAP1
Full name:	DNA methyltransferase 1 associated protein 1
Synonyms:	EAF2; SWC4; MEAF2; DNMAP1; DNMTAP1
SwissProt:	Q9NPF5
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human gastric cancer; Human thyroid cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	53 kDa
WB Positive control:	231 and Hela cell lysates
WB Recommended dilution:	200-1000



