

兔抗 EIF4A2 多克隆抗体

中文名称： 兔抗 EIF4A2 多克隆抗体

英文名称： Anti-EIF4A2 rabbit polyclonal antibody

别 名： eukaryotic translation initiation factor 4A2; DDX2B; EIF4A; EIF4F; BM-010; EIF4A-II; eIF-4A-II

相关类别： 一抗

储 存： 冷冻 (-20°C)

宿 主： Rabbit

抗 原： EIF4A2

反应种属： Human, Mouse, Rat

标 记 物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.
Applications:	ELISA, IHC
Name of antibody:	EIF4A2
Immunogen:	Fusion protein of human EIF4A2

Full name:	eukaryotic translation initiation factor 4A2
Synonyms:	DDX2B; EIF4A; EIF4F; BM-010; eIF4A-II; eIF-4A-II
SwissProt:	Q14240
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and Human esophagus cancer
IHC Recommend dilution:	50-200



