

全国订货电话 4008-723-722

兔抗 MRPL10 多克隆抗体

- 中文名称: 兔抗 MRPL10 多克隆抗体
- 英文名称: Anti-MRPL10 rabbit polyclonal antibody
- 别名: L10MT; MRPL8; RPML8; MRP-L8; MRP-L10
- 储存: 冷冻(-20℃)避光
- 抗原: MRPL10
- 宿 主: Rabbit
- 反应种属: Human
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

	Mammalian mitachandrial vibacamal protains are anaded
Background:	Mammalian mitochondrial ribosomal proteins are encoded
	by nuclear genes and help in protein synthesis within the
	mitochondrion. Mitochondrial ribosomes (mitoribosomes) c
	onsist of a small 28S subunit and a large 39S subunit. The
	y have an estimated 75% protein to rRNA composition co
	mpared to prokaryotic ribosomes, where this ratio is revers
	ed. Another difference between mammalian mitoribosomes
	and prokaryotic ribosomes is that the latter contain a 5S r $% \left({{\left[{{n_{\rm{s}}} \right]}} \right)$
	RNA. Among different species, the proteins comprising the
	mitoribosome differ greatly in sequence, and sometimes in
	biochemical properties, which prevents easy recognition by
	sequence homology. This gene encodes a 39S subunit prot
	ein. Sequence analysis identified three transcript variants th
	at encode two different isoforms. A pseudogene correspon





	ding to this gene is found on chromosome 5q.
Applications:	WB
Name of antibody:	MRPL10
Immunogen:	Synthesized peptide derived from internal of human MRPL 10.
Full name:	mitochondrial ribosomal protein L10
Synonyms :	L10MT; MRPL8; RPML8; MRP-L8; MRP-L10
SwissProt:	Q7Z7H8
WB Predicted band size:	29 kDa
WB Positive control:	HepG2 cells, COLO cells and 293 cells lysates
WB Recommended dilution:	500-3000

