

兔抗 RPL10A 多克隆抗体

中文名称：兔抗 RPL10A 多克隆抗体

英文名称： Anti-RPL10A rabbit polyclonal antibody

别名： ribosomal protein L10a; L10A; CSA19; NEDD6; Csa-19

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： RPL10A

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L1P family of ribosomal proteins. It is located in the cytoplasm. The expression of this gene is downregulated in the thymus by cyclosporin-A (CsA), an immunosuppressive drug. Studies in mice have shown that the expression of the ribosomal protein L10a gene is downregulated in neural precursor cells during development. This gene previously was referred to as NEDD6 (neural prec

	ursor cell expressed, developmentally downregulated 6), but it has been renamed RPL10A (ribosomal protein 10 a). As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.
Applications:	ELISA, WB, IHC
Name of antibody:	RPL10A
Immunogen:	Fusion protein of human RPL10A
Full name:	ribosomal protein L10a
Synonyms:	L10A; CSA19; NEDD6; Csa-19
SwissProt:	P62906
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer and Human thyroid cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	25 kDa
WB Positive control:	Rat liver tissue, Mouse kidney tissue, Mouse lung tissue , HUVEC, Jurkat, HeLa, HepG2 cell lysates
WB Recommended dilution:	500-2000



