

兔抗 SLC22A12 多克隆抗体

| | |
|-------|---|
| 中文名称: | 兔抗 SLC22A12 多克隆抗体 |
| 英文名称: | Anti-SLC22A12 rabbit polyclonal antibody |
| 别名: | solute carrier family 22 member 12; RST; OAT4L; URAT1 |
| 相关类别: | 一抗 |
| 储存: | 冷冻 (-20℃) |
| 宿主: | Rabbit |
| 抗原: | SLC22A12 |
| 反应种属: | Human |
| 标记物: | Unconjugate |
| 克隆类型: | rabbit polyclonal |

技术规格

| | |
|--------------------------|--|
| Background: | The protein encoded by this gene is a member of the organic anion transporter (OAT) family, and it acts as a urate transporter to regulate urate levels in blood. This protein is an integral membrane protein primarily found in epithelial cells of the proximal tubule of the kidney. An elevated level of serum urate, hyperuricemia, is associated with increased incidences of gout, and mutations in this gene cause renal hypouricemia type 1. Alternative splicing results in multiple transcript variants. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | SLC22A12 |
| Immunogen: | Synthetic peptide of human SLC22A12 |
| Full name: | solute carrier family 22 member 12 |
| Synonyms: | RST; OAT4L; URAT1 |

| | |
|------------------------------------|---|
| SwissProt: | Q96S37 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human cervical cancer and Human colorectal cancer |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 60 kDa |
| WB Positive control: | 293T cell lysate |
| WB Recommended dilution: | 500-2000 |



