

兔抗 PSPC1 多克隆抗体

- 中文名称: 兔抗 PSPC1 多克隆抗体
- 英文名称: Anti-PSPC1 rabbit polyclonal antibody
- 别 名: paraspeckle component 1; PSP1
- 抗 原: PSPC1
- 储 存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human, Mouse, Rat
- 标 记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

| Background: | This gene encodes a nucleolar protein that localizes to p unctate subnuclear structures that occur close to splicing speckles, known as paraspeckles. These paraspeckles are composed of RNA-protein structures that include a non- coding RNA, NEAT1/Men epsilon/beta, and the Drosophil a Behavior Human Splicing family of proteins, which incl ude the product of this gene and the P54NRB/NONO an d PSF/SFPQ proteins. Paraspeckles may function in the c ontrol of gene expression via an RNA nuclear retention mechanism. The protein encoded by this gene is found i n paraspeckles in transcriptionally active cells, but it local izes to unique cap structures at the nucleolar periphery |
|-------------|---|
| | when RNA polymerase II transcription is inhibited, or dur |



全国订货电话 4008-723-722

| | ing telophase. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene, which is also located on chromosome 13, has been identified. |
|-----------------------------|--|
| Applications: | ELISA, IHC |
| Name of antibody: | PSPC1 |
| Immunogen: | Fusion protein of human PSPC1 |
| Full name: | paraspeckle component 1 |
| Synonyms: | PSP1 |
| SwissProt: | Q8WXF1 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human thyroid cancer |
| IHC Recommend dilution: | 30-150 |





全国订货电话 4008-723-722

