

兔抗 SLC27A4 多克隆抗体

中文名称：兔抗 SLC27A4 多克隆抗体

英文名称：Anti-SLC27A4 rabbit polyclonal antibody

别名：IPS; FATP4; ACSVL4

抗原：SLC27A4

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes a member of a family of fatty acid transport proteins, which are involved in translocation of long-chain fatty acids across the plasma membrane. This protein is expressed at high levels on the apical side of mature enterocytes in the small intestine, and appears to be the principal fatty acid transporter in enterocytes. Clinical studies suggest this gene as a candidate gene for the insulin resistance syndrome. Mutations in this gene have been associated with ichthyosis pr

| | |
|------------------------------------|---|
| | ematurity syndrome. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | SLC27A4 |
| Immunogen: | Fusion protein of human SLC27A4 |
| Full name: | solute carrier family 27 (fatty acid transporter), member 4 |
| Synonyms: | IPS; FATP4; ACSVL4 |
| SwissProt: | Q6P1M0 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human brain and human esophagus cancer |
| IHC Recommend dilution: | 25-100 |
| WB Predicted band size: | 72 kDa |
| WB Positive control: | PC-3 , Hela , 293T and HEPG2 cell lysates |
| WB Recommended dilution: | 200-1000 |





